arXiv:hep-ph/0408015v3 13 Oct 2005

IFUP-TH/27-2004
IFT-2004/21
ECT*-04-09
hep-ph/0408015

Low-scale standard supersymmetric leptogenesis

Martti Raidal[1], **Alessandro Strumia**[2] and **Krzysztof Turzyński**[3]

[1] *National Institute of Chemical Physics and Biophysics, Tallinn 10143, Estonia*
[2] *Dipartimento di Fisica dell'Università di Pisa and INFN, Italia*
[3] *Institute of Theoretical Physics, Warsaw University, ul. Hoża 69, 00-681 Warsaw, Poland*

Abstract

Strictly adhering to the standard supersymmetric seesaw mechanism, we present a neutrino mass model which allows successful standard thermal leptogenesis compatible with gravitino cosmology. Some neutrino Yukawa couplings are naturally much larger than the naive estimates following from the seesaw formula. This leads to large BR($\mu \to e\gamma$), detectable in the next round of experiments. Ratios of $\mu \to e\gamma$, $\tau \to e\gamma$ and $\tau \to \mu\gamma$ branching ratios are predicted in terms of the measurable neutrino mass matrix.

1 Introduction

The observed neutrino masses can be explained by adding to the Standard Model (SM) one heavy right-handed neutrino N_i per generation [1]:

$$\mathscr{L} = \mathscr{L}_{\rm SM} + \bar{N}_i i\partial\!\!\!/ N_i + \left(\boldsymbol{\lambda}_{i\ell} N_i L_\ell H + \frac{1}{2}\boldsymbol{M}_{ij} N_i N_j + \text{h.c.}\right). \tag{1}$$

The neutrino mass matrix depends on the unknown right-handed neutrino mass matrix $\boldsymbol{M}$ (with eigenvalues $M_1 < M_2 < M_3$) and neutrino Yukawa couplings $\boldsymbol{\lambda}$ as

$$\boldsymbol{m}_\nu = -v^2 \boldsymbol{\lambda}^T \boldsymbol{M}^{-1} \boldsymbol{\lambda}, \tag{2}$$

suggesting that the masses of the right-handed neutrinos are below $10^{14}\,\text{GeV}$. Unlike the SM, this extension can generate the observed baryon asymmetry

$$\frac{n_B}{n_\gamma} = (6.15 \pm 0.25) \times 10^{-10} \tag{3}$$

via thermal leptogenesis in out-of-equilibrium decays of the lightest right-handed neutrino [2].

Various considerations motivate supersymmetry broken by Fermi-scale soft terms. Supersymmetric seesaw models have new signals: quantum effects imprint the lepton flavour violation of the neutrino Yukawa couplings in the left-handed slepton masses, giving rise to potentially observable $\mu \to e\gamma$ and related lepton-flavour violating (LFV) processes. Their rates depend on the unknown (but measurable) sparticle spectrum and, more importantly, on the unknown neutrino Yukawa couplings $\boldsymbol{\lambda}$. Since neutrino masses tell us only $\boldsymbol{\lambda}^T \boldsymbol{M}^{-1} \boldsymbol{\lambda}$, LFV rates cannot be predicted in terms of low-energy observables.

In addition, standard supersymmetric leptogenesis faces a potential problem: gravitino decays destroy Big Bang Nucleosynthesis (BBN), unless the maximal 'reheating' temperature of the hot Big Bang $T_{\rm RH}$ is less than about $10^7\,\text{GeV}$ (the precise value depends on the gravitino mass and on its hadronic branching ratio) [3].[1] On the other hand, successful minimal leptogenesis requires [15, 17, 18]

$$T_{\rm RH} > 2 \times 10^9\,\text{GeV}, \tag{4}$$

if zero initial abundance of right-handed neutrinos is assumed and only the leading $\mathcal{O}(M_1/M_{2,3})$ terms are included in the calculation of the CP asymmetry ε_1 in decays $N_1 \to LH, \bar{L}H^*$ of the lightest right-handed neutrino N_1. These terms originate from the same effective operator $(LH)^2$ which also gives the contributions $\tilde{m}_{2,3}$ of the heavier $N_{2,3}$ to neutrino masses[2]. Thanks to this close relation, the naive estimate $\varepsilon_1 \sim 3\tilde{m}_{2,3}M_1/8\pi v^2$ can be improved to the rigorous Davidson-Ibarra bound [17, 21]

---

[1] This bound does not apply if gravitinos are the lightest stable SUSY particles [4]. However, in such a case, BBN can be destroyed by decays of the next-to-lightest SUSY particles [3]. Other scenarios of leptogenesis that avoid the gravitino problem, e.g. resonant leptogenesis with almost degenerate singlet neutrino masses [5, 6, 7, 8, 9], soft leptogenesis [10, 11], non-thermal leptogenesis [12], and leptogenesis with N_1 or $\tilde{N}_1$-dominated universe [13, 14, 15] have been proposed. The gravitino problem is absent in models where the gravitino is ultraheavy or not present at all [16].

$$|\varepsilon_1| \le \varepsilon_1^{\rm DI} = \frac{3}{8\pi}\frac{(m_3 - m_1)M_1}{v^2} \tag{5}$$

where $m_1 < m_2 < m_3$ are light neutrino masses. Combined with precise studies of the dynamics of thermal leptogenesis [15] this implies eq. (4).

Higher orders corrections to this formula, suppressed by powers of $M_1/M_{2,3}$, are not directly related to neutrino masses and give the additional contribution

$$\delta\varepsilon_1 \simeq \frac{3}{16\pi}\frac{\tilde{m}_{2,3}M_1}{v^2}\left(\frac{M_1}{M_{2,3}}\right)^2 . \tag{6}$$

A CP asymmetry much above the bound (5) is obtained for $\tilde{m}_{2,3}/m_{2,3} \gg (M_{2,3}/M_1)^2$ [20]. In other words, one needs that the N_2 and N_3 contributions to neutrino masses $\tilde{m}_{2,3}$ are orders of magnitude larger than the neutrino masses $m_{2,3}$ themselves, i.e. these contributions have to cancel each other. Barring anthrophic arguments and in absence of an underlying theoretical justification this would be an unlikely fine-tuning. While both papers in Ref. [20] claim that it seems possible to justify such cancellations, the suggested pseudo-Dirac structure fails because it controls neutrino masses, but at the price of suppressing the CP-asymmetry of eq. (6) [20].

The goal of this paper consists in finding the class of models which realizes the above mechanism naturally and in studying their implications for the rates of LFV processes. In section 2 we present the texture of the right-handed neutrinos mass matrix needed to control the appearance of the Yukawa couplings in the light neutrino mass matrix so that these cancellations occur naturally, and outline how it can be justified by flavour models. We also address the issue of the size of effective neutrino mass $\tilde{m}_1$ and show that it can be naturally small as required by successful leptogenesis. In section 3 we perform a phenomenological study of such models relating the model parameters to the neutrino oscillation measurements. We show that the proposed class of models predicts ratios of $\mu \to e\gamma$, $\tau \to e\gamma$ and $\tau \to \mu\gamma$ branching ratios in terms of the measurable neutrino mass matrix. In section 4 we show that such LFV rates must be sizable in the region of the parameter space that allows successful thermal leptogenesis compatibly with the gravitino constraint. In section 5 we discuss the stability of predictions with respect to small perturbations from the 'exact' texture. Results are summarized in section 6.

---

[2] In this limit the two 'vertex' and 'self-energy' loop diagrams collapse into a single diagram. This observation could have allowed avoiding past debates (solved by [19]) on whether the 'self-energy' diagram contributes to ε_1. See [20] for a quantitative definition and discussion of $\tilde{m}_{2,3}$.

2 Building models with $|\varepsilon_1| \gg \varepsilon_1^{\rm DI}$

In this section we outline the basic features of our neutrino mass model which naturally accommodates an enhanced CP-asymmetry $|\delta\varepsilon_1| \gg \varepsilon_1^{\rm DI}$. For later convenience we introduce an unusual notation, writing the neutrino Yukawa matrix as

$$\boldsymbol{\lambda}_{i\ell} = \begin{pmatrix} \lambda_1^e & \lambda_1^\mu & \lambda_1^\tau \\ \lambda_2^e & \lambda_2^\mu & \lambda_2^\tau \\ \lambda_3^e & \lambda_3^\mu & \lambda_3^\tau \end{pmatrix} = \begin{pmatrix} \vec{\lambda}_1 \\ \vec{\lambda}_2 \\ \vec{\lambda}_3 \end{pmatrix} \tag{7}$$

where $\vec{\lambda}_{1,2,3}$ are 3-vectors with $\ell = \{e, \mu, \tau\}$ components. We assume that the entries orders of magnitude larger than what naïvely suggested by the seesaw (needed to get large $\tilde{m}_{2,3}$) are all contained in the Yukawa couplings $\vec{\lambda}_3$ of N_3. For the moment we assume the simplest hypothesis suggested by data: 'lopsided' Yukawa couplings [22], i.e. that each $\vec{\lambda}_i$ has comparable entries $\lambda_i^{e,\mu,\tau}$, which naturally leads to large mixing angles.[3] Our assumption therefore reads

$$\lambda_3^\ell \gg \lambda_{1,2}^\ell. \tag{8}$$

Finally, our key assumption is that, in the basis where eq. (8) holds, the right-handed neutrino mass matrix has the following form

$$\boldsymbol{M} = \begin{array}{c} \\ N_1 \\ N_2 \\ N_3 \end{array} \begin{array}{c} \begin{array}{ccc} N_1 & N_2 & N_3 \end{array} \\ \begin{pmatrix} M_{11} & 0 & 0 \\ 0 & 0 & M_{23} \\ 0 & M_{23} & M_{33} \end{pmatrix} \end{array}. \tag{9}$$

The assumption $\boldsymbol{M}_{12} = \boldsymbol{M}_{22} = 0$ implies vanishing 33 and 13 entries in the inverse mass matrix

$$\boldsymbol{M}^{-1} = \begin{pmatrix} M_{11}^{-1} & 0 & 0 \\ 0 & -M_{33}/M_{23}^2 & M_{23}^{-1} \\ 0 & M_{23}^{-1} & 0 \end{pmatrix} \tag{10}$$

that enters the seesaw formula

$$\boldsymbol{m}_\nu = -v^2 \sum_{i,j} (\boldsymbol{M}^{-1})_{ij}\, \vec{\lambda}_i \otimes \vec{\lambda}_j. \tag{11}$$

Therefore, the large Yukawa couplings $\vec{\lambda}_3$ do not generate large contributions to the neutrino masses of order $\lambda_3^2 v^2/M_{33}$. In the right-handed neutrino mass basis, both $N_2^{\rm mass}$ and $N_3^{\rm mass}$ give large contributions $\tilde{m}_{2,3} \sim \lambda_3^2 v^2/M_{2,3}$ to neutrino masses, but they naturally cancel out.

The matrix $\boldsymbol{M}^{-3}$, which controls the 'higher order' contributions to ε_1, has no special structure in the N_2, N_3 sector:

$$\boldsymbol{M}^{-3} = \begin{pmatrix} M_{11}^{-3} & 0 & 0 \\ 0 & \mathcal{O}(M_{23}^{-3}) & \mathcal{O}(M_{23}^{-3}) \\ 0 & \mathcal{O}(M_{23}^{-3}) & -M_{33}/M_{23}^4 \end{pmatrix} \tag{12}$$

---

[3] *A posteriori* in section 3 we will find that for $\tilde{m}_1 \ll m_{\rm sun}$ the model is predictive enough that our *a priori* assumption of lopsided $\vec{\lambda}_{2,3}$ is required by observed neutrino masses and mixings.

and its 33 entry gives the dominant contribution to ε_1, proportional to λ_3^2.

A crucial issue for successful leptogenesis is the smallness of the effective neutrino mass $\tilde{m}_1$, needed to have enough out-of-equilibrium N_1 decays. The further assumption $\boldsymbol{M}_{13} = 0$ (together with $\boldsymbol{M}_{12} = 0$) guarantees that the large Yukawa coupling λ_3 does not contribute to $\tilde{m}_1$. Indeed, N_1 is already the lightest mass-eigenstate which operates leptogenesis, with mass $M_1 = M_{11}$ and $\tilde{m}_1 = (\vec{\lambda}_1^* \cdot \vec{\lambda}_1)v^2/M_{11}$, which is therefore small.

Thus the texture described between equations (7) and (9) has, in principle, the features which allow successful leptogenesis and reproducing the observed neutrino masses and mixings. We shall discuss these issues more thoroughly in the following sections.

So far we identified a special basis in which our initial problem gets reduced to the simpler problem of explaining why certain entries are small. We now discuss how the needed texture can be theoretically justified by building appropriate flavour models. This second step usually does not lead to further insights, because usually there are many possible ways of justifying small couplings. This will turn out to be true also in our case, which however presents a potential difficulty: the texture eq. (9) need strict zeroes in specific entries. Therefore we here explicitly address only this potentially difficult issue, briefly presenting one example with $M_{23} \gtrsim M_{33}$ (models with $M_{23} \lesssim M_{33}$ look more complicated). We introduce a U(1) flavour symmetry under which N_1, N_2, N_3, $(L_{e,\mu,\tau}H)$ have charges $0, 1, -1, 1$, respectively, and which is broken by the vev of a flavon superfield φ_+ with charge 1. This implies $\boldsymbol{M}_{12} = \boldsymbol{M}_{22} = 0$, $\vec{\lambda}_1 = \vec{\lambda}_2 = 0$ and allows a 'lopsided' $\vec{\lambda}_3$. One can force $\boldsymbol{M}_{13} = 0$ by assuming that φ_+ has charge $2/3$ (rather than 1), or by adding extra symmetries that act only on N_1 or by assuming that N_1 and $N_{2,3}$ have wave-functions localized in extra dimensions and with small overlap among them. Finally, assuming that the flavon φ_- with charge equal to $\varphi_+^\dagger$ has a small vev $\langle\varphi_-\rangle \ll \langle\varphi_+\rangle$, which is natural in supersymmetric models[4], allows small 'lopsided' $\vec{\lambda}_{1,2}$ and small $\boldsymbol{M}_{12}$, $\boldsymbol{M}_{22}$. Note that the mechanism that ensures $\boldsymbol{M}_{13} = 0$ also breaks the relation $\lambda_1^2 \sim \lambda_2$, expected from our minimal charge assignment. One possibility of generating λ_1 and λ_2 in the phenomenologically interesting range (discussed later) is the introduction of four flavons with charges $\pm 2/3$ and ± 1 and with appropriate vevs[5]. The U(1) flavour symmetry can be extended to the quarks and right-handed leptons in a SU(5)-invariant way, reproducing the masses and mixings of all SM fermions. It should be clear that proceeding as outlined above we could build many full models. We here prefer to study the general testable consequences.

Small entries of $\boldsymbol{M}$ and $\boldsymbol{\lambda}$ are stable under quantum corrections. This can be easily seen in SUSY models, where quantum corrections only renormalize kinetic terms. Furthermore, at tree level, $N_{1,2,3}$ in general already have a non-canonical kinetic matrix. Since it does not affect the seesaw mechanism we can ignore it.

---

[4] The F-term potential, being invariant under the complexified global flavour symmetry, determines at most the product $\langle\varphi_+\varphi_-\rangle$. The individual vevs might be fixed by SUSY-breaking corrections; the desired hierarchy $\langle\varphi_-\rangle \ll \langle\varphi_+\rangle$ can be naturally obtained e.g. when $\langle\varphi_+\varphi_-\rangle = 0$ in the SUSY-limit. 'Supersymmetric zeroes' are discussed in [23].

[5] We thank S. Davidson for suggesting models with such kinds of charge assignments and for indicating us the features we presented.

3 Flavour violation

In this section we perform a thorough phenomenological study of the proposed scenario with the aim to relate model parameters to the neutrino measurements. We start with studying the values of the neutrino Yukawa couplings $\vec{\lambda}_{1,2,3}$ with emphasis on the entries of the largest Yukawa couplings $\vec{\lambda}_3$, which are the dominant source of the LFV effects. So far we assumed that each λ_i is roughly equally distributed among flavours. We shall discuss carefully, how much this assumption is required by the observed large atmospheric and solar mixing angles, obtaining precise predictions for the flavour content of the neutrino Yukawa couplings.

We choose $\tilde{m}_1 \sim 10^{-3}\,\mathrm{eV}$, which yields

$$\lambda_1 \equiv (\vec{\lambda}_1^* \cdot \vec{\lambda}_1)^{1/2} = \frac{\sqrt{\tilde{m}_1 M_1}}{v} \sim 10^{-6} \frac{M_1}{10^6\,\mathrm{GeV}}. \tag{13}$$

Together with large λ_3, these choices make n_B as large as possible[6]. Then the neutrino mass matrix is dominated by the term

$$\boldsymbol{m}_\nu = -\frac{v^2}{M_{23}} \left(\vec{\lambda}_3 \otimes \vec{\lambda}_2 + \vec{\lambda}_2 \otimes \vec{\lambda}_3 \right), \tag{14}$$

which can give both the atmospheric and solar masses $m_2 = \sqrt{\Delta m^2_{\rm sun}}$, $m_3 = \sqrt{\Delta m^2_{\rm atm}}$ for

$$\lambda_2 \sim \frac{m_{2,3} M_{2,3}}{\lambda_3 v^2} \sim 10^{-8} \frac{1}{\lambda_3} \frac{M_{2,3}}{10^7\,\mathrm{GeV}}. \tag{15}$$

We now invert the relation (14) between seesaw parameters and observable neutrino masses and mixings. The analytic computation will be performed in an unusual but convenient way: as suggested by the vector notation $\vec{\lambda}_i$ we will use vector calculus.[7]

---

[6] This might be however unnecessary, allowing smaller or larger values of λ_1 and smaller values of λ_3. We will discuss this issue later.

[7] All our subsequent expressions can be converted into traditional notations by replacing vectors with their components. Models with neutrino masses and/or leptogenesis dominated by two right-handed neutrinos have been studied previously in the literature, but only performing generic analyses [24, 25] or focusing on specific models [27] different from the ones under consideration.

The computation below could be alternatively performed in a more indirect way using the R-matrix introduced in [28], after finding the specific class of R matrices corresponding to our class of models. Such R matrices can be parametrized as

$$R \simeq \begin{pmatrix} 1 & 0 & 0 \\ 0 & z & iz \\ 0 & -iz & z \end{pmatrix} \qquad \text{for } |z| \gg 1.$$

The phase of the complex number z does not play any role. See also [20, 25]. In particular, our predictions for LFV rates are equivalent to appropriate limits of equations (18), (19) in [25].

Computing neutrino masses $m_i \geq 0$ and neutrino mass eigenstates $\hat{\nu}_i$ amounts to decompose the neutrino mass matrix given in eq. (14) as

$$\boldsymbol{m}_\nu = \sum_{i=1}^{3} m_i \, \hat{\nu}_i^* \otimes \hat{\nu}_i^* \tag{16}$$

where $\hat{\nu}_i$ are orthonormal vectors spanning the flavour space of the left-handed doublets. In the basis of charged lepton mass eigenstates they are usually parametrized as

$$\hat{\nu}_3^\ell = \boldsymbol{V}_\nu^{\ell 3} \qquad \hat{\nu}_2^\ell = \boldsymbol{V}_\nu^{\ell 2} e^{i\alpha} \qquad \hat{\nu}_1^\ell = \boldsymbol{V}_\nu^{\ell 1} e^{i\beta} \tag{17}$$

where $\boldsymbol{V}_\nu$ is the CKM-like part of the neutrino mixing matrix and α, β are Majorana phases.

Defining the moduli $\lambda_i \equiv (\vec{\lambda}_i^* \cdot \vec{\lambda}_i)^{1/2}$, the unit versors $\hat{\lambda}_i \equiv \vec{\lambda}_i / \lambda_i$ and $\vec{\lambda}_3^* \cdot \vec{\lambda}_2 \equiv \lambda_2 \lambda_3 c e^{2i\psi}$, where $0 \leq c \leq 1$ is the 'cosine' of the angle between $\vec{\lambda}_2$ and $\vec{\lambda}_3$, we get:

$$m_1 = 0, \qquad m_{2,3} = \lambda_2 \lambda_3 (1 \mp c) v^2 / M_{23}. \tag{18}$$

The observed moderate mass ratio $m_3/m_2 \sim 5$ between the atmospheric and solar masses is reproduced for $c \approx \cos 45^o$, which does not require any special alignment of $\hat{\lambda}_2$ and $\hat{\lambda}_3$ in the flavour space, consistently with our assumption of 'lopsided' vectors $\vec{\lambda}_{2,3}$. In contrast, the inverted hierarchy of the neutrino masses requires $c \simeq \Delta m^2_{\rm sun}/4\Delta m^2_{\rm atm} \sim 10^{-2}$, which means that $\hat{\lambda}_2$ and $\hat{\lambda}_3$ have to be almost orthogonal in the flavour space. Such an alignment would clearly be less natural and need further justification.[8]

The neutrino mass eigenstates are

$$\hat{\nu}_1 \propto \hat{\lambda}_3 \times \hat{\lambda}_2 \qquad \hat{\nu}_2 = \frac{-e^{i\psi}\hat{\lambda}_3^* + e^{-i\psi}\hat{\lambda}_2^*}{\sqrt{2(1-c)}} \qquad \hat{\nu}_3 = i\frac{e^{i\psi}\hat{\lambda}_3^* + e^{-i\psi}\hat{\lambda}_2^*}{\sqrt{2(1+c)}}. \tag{19}$$

Inverting this relation we get

$$\hat{\lambda}_{3,2} = e^{\pm i(\psi \pm \pi/2)} \frac{\hat{\nu}_3^* \pm i\sqrt{m_2/m_3}\,\hat{\nu}_2^*}{\sqrt{1 + m_2/m_3}} \tag{20}$$

in terms of the low energy observables (not affected by the phase ψ). Coming back to the traditional notation the flavour content of the big neutrino Yukawa coupling is

$$\hat{\lambda}_3^\ell = i e^{i\psi} \frac{\boldsymbol{V}_\nu^{\ell 3} + i e^{i\alpha} \boldsymbol{V}_\nu^{\ell 2} \sqrt{m_2/m_3}}{\sqrt{1 + m_2/m_3}}. \tag{21}$$

---

[8] If nevertheless neutrinos turn out to have an inverted spectrum, our predictions for this case can be found by substituting $V_\nu^{\ell 3}$ with $V_\nu^{\ell 1}$ in eq. (21).

Changing α by π is equivalent to swapping $\hat{\lambda}_3$ and $\hat{\lambda}_2$. In view of observed large neutrino mixing angles, the Yukawa coupling vectors $\hat{\lambda}_2^\ell$ and $\hat{\lambda}_3^\ell$ turn out to generically have 'lopsided' form, which justifies our heuristic assumption made in Section 2.[9]

It follows from eq. (21) that the model predicts

$$\frac{\mathrm{BR}(\tau \to \mu\gamma)}{\mathrm{BR}(\mu \to e\gamma)} = b\frac{|\lambda_3^\tau|^2}{|\lambda_3^e|^2}, \qquad \frac{\mathrm{BR}(\tau \to e\gamma)}{\mathrm{BR}(\mu \to e\gamma)} = b\frac{|\lambda_3^\tau|^2}{|\lambda_3^\mu|^2} \tag{22}$$

(where $b = BR(\tau \to \mu\bar{\nu}_\mu\nu_\tau) = 0.18$) in terms of the neutrino masses, mixings and CP phases measurable at low energy. We fix the already measured quantities to their present best-fit values

$$\theta_{23} = \frac{\pi}{4}, \qquad \tan^2\theta_{12} = 0.4, \qquad \frac{m_3}{m_2} = 5, \tag{23}$$

and show in fig. 1a the predicted range of ratios among different branching ratios. We fix θ_{13} at a few possible values and freely vary the remaining unmeasured parameters, i.e. the Majorana phase α and the CP-phase ϕ in neutrino oscillations. The dashed line is an example of the predicted range if all parameters except α were measured. The model predicts that $\mu \to e\gamma$, $\tau \to e\gamma$ and $\tau \to \mu\gamma$ have comparable branching ratios. Since $\mathrm{BR}(\mu \to e\gamma)$ is presently the most strongly constrained, only $\mu \to e\gamma$ seems observable in the next round of experiments.

Note that it is not possible to have $\lambda_3^e = 0$ and therefore negligibly small $\mathrm{BR}(\mu \to e\gamma)$ and $\mathrm{BR}(\tau \to e\gamma)$, since this would require

$$\tan\theta_{13} = \sqrt{m_2/m_3}\sin\theta_{12} \tag{24}$$

which violates the 'Chooz' [26] constraint on θ_{13}. More precisely, taking into account that θ_{13} is also constrained by 'solar' and 'atmospheric' experiments, by performing a global fit we find $\sin^2\theta_{13} = 0.007 \pm 0.016$: data disfavour the relation (24) at about 3σ.

As usual, the predicted LFV rates depend on sparticle masses which can be measured at colliders. Taking into account naturalness considerations and experimental bounds and hints, we give our numerical examples for $m_0 = 100\,\mathrm{GeV}$, $M_{1/2} = 200\,\mathrm{GeV}$, $A_0 = 0$, $\tan\beta = 5$. $\mathrm{BR}(\mu \to e\gamma)$ is then approximately given by

$$\mathrm{BR}(\mu \to e\gamma) \approx 2 \cdot 10^{-7} r |\boldsymbol{Y}_{e\mu}|^2 \qquad \text{where} \qquad \boldsymbol{Y} = \boldsymbol{\lambda}^\dagger \ln\left(\frac{M_{\mathrm{GUT}}^2}{\boldsymbol{M}\boldsymbol{M}^\dagger}\right)\boldsymbol{\lambda} \tag{25}$$

and $r \approx (\tan\beta/5)^2(200\,\mathrm{GeV}/M_{\mathrm{SUSY}})^4$ equals 1 at our reference point.[10] Figures 1 and 2 show results of a precise computation, performed by solving numerically the full set of

---

[9] Only if future experiments establish θ_{13} much below present bounds, eq. (19) will need unnatural cancellations indicating the necessity of a flavour symmetry more complicated than 'lopsided'. Our predictions would remain unchanged, as they follow from the texture in eq. (9), not from the flavour models that can justify it.

[10] See, however, [25] for a discussion of potentially sizable corrections to this simple estimate. Furthermore, if M_1 is lowered down to Fermi-scale energies tree-level exchange of right-handed neutrinos gives extra contributions to LFV rates beyond the supersymmetric loop contributions discussed here.



Figure 1: *a) Ratios of LFV decays allowed for* $\theta_{13} = 0$ *(thick black line),* $0.05, 0.1, 0.15$ *for arbitrary values of the Majorana phase* α *and of the CP- phase* ϕ *in neutrino oscillations. The dashed line shows the residual allowed range assuming that future measurements will fix* $\theta_{13} = 0$ *and* $\phi = 1$*. b) Range of* $\mathrm{BR}(\mu \to e\gamma)$ *for arbitrary values of the Majorana phase* α *as a function of the CP Dirac phase* ϕ *for different values of* θ_{13}*. The factor* $r \approx (\tan\beta/5)^2(200\,\mathrm{GeV}/M_{\mathrm{SUSY}})^4$ *equals one at our reference SUSY spectrum. We fixed* $\lambda_3 = 0.1$*,* $M_1 = 10^6\,\mathrm{GeV}$*,* $M_3/M_2 = 2$*.*

the renormalization group equations of the supersymmetric seesaw model (see e.g. [9, 29]) and using exact formulae of [30].

In fig. 1b we show how $\mathrm{BR}(\mu \to e\gamma)$ depends on θ_{13}, ϕ and α at fixed values of high-energy parameters. In fig. 2a we show how $\mathrm{BR}(\mu \to e\gamma)$ (for $r = 1$, $\theta_{13} = 0.1$, $\phi = 1$ and minimized with respect to Majorana phases) depends on the main high-energy parameter, λ_3. As illustrated in the figure and discussed in the next section, successful thermal leptogenesis demands a value of λ_3 that implies $\mu \to e\gamma$ at a level observable in the next round of experiments (unless sparticles are much above the Fermi scale and unless SUSY breaking is mediated to sleptons at energies below M_1).

4 Leptogenesis

Gravitino cosmology probably requires M_1 a few orders of magnitude below its minimal value $M_1^{\mathrm{DI}} \sim 2 \times 10^9\,\mathrm{GeV}$ allowed by the bound (5) on the CP asymmetry ε_1 [3, 17]. In



Figure 2: *As function of* λ_3 *and of* M_2/M_1 *we plot contour lines of: (a) the minimal value of* $\mathrm{BR}(\mu \to e\gamma)$*; the factor* $r \approx (\tan\beta/5)^2(200\,\mathrm{GeV}/M_{\mathrm{SUSY}})^4$ *equals one at our reference SUSY spectrum. (b) the maximal baryon asymmetry* n_B/n_γ*. The shaded region covers the allowed region. We assumed* $\theta_{13} = 0.1$, $\phi = 1$, $M_1 = 10^6\,\mathrm{GeV}$, $M_3/M_2 = 2$, $\tilde{m}_1 = 10^{-3}\,\mathrm{eV}$, *but the main conclusions do not depend on these choices, as explained in the text.*

our model this can be achieved thanks to a ε_1 orders of magnitude larger than $\varepsilon_1^{\mathrm{DI}}$:

$$|\varepsilon_1| \leq \varepsilon_1^{\mathrm{DI}} + \frac{3}{16\pi}\frac{(M_3 - M_2)M_1^3}{M_2^2 M_3^2}\lambda_3^2. \tag{26}$$

(In the limit $M_3 \gg M_2$ the maximal CP-asymmetry reduces to the estimate of [20]). The CP asymmetry needed to reach a given $M_1 < M_1^{\mathrm{DI}}$ is $|\varepsilon_1| = \varepsilon_1^{\mathrm{DI}} M_1/M_1^{\mathrm{DI}}$ (in the absence of additional washout effects) and is obtained for

$$\lambda_3 \approx \frac{\sqrt{m_3 M_1^{\mathrm{DI}}}}{v}\left(\frac{M_{2,3}}{M_1}\right)^{3/2} \approx 0.1\left(\frac{M_{2,3}}{10\,M_1}\right)^{3/2}, \tag{27}$$

where we approximated $M_{2,3} \approx M_2 \approx M_3$. Relatively large values of λ_3 are needed irrespectively of the value of M_1/M_1^{DI}, since for smaller M_1 the asymmetry ε_1 needs more enhancement by the Yukawa couplings, whose naïve values become smaller. The CP asymmetry is saturated for $|\mathrm{Im}\,\hat{\lambda}_1 \cdot \hat{\lambda}_3^*| = 1$, which can be achieved compatibly with all experimental constraints on neutrino masses and mixings. This becomes particularly clear for $\tilde{m}_1 \ll m_{\mathrm{sun}}$, because in this case $\vec{\lambda}_1$ negligibly contributes to the neutrino mass matrix, so that $\hat{\lambda}_1$ remains fully undetermined. 'Lopsided' Yukawa couplings suggest the generic expectation $|\mathrm{Im}\,\hat{\lambda}_1 \cdot \hat{\lambda}_3^*| \lesssim 1$.

The ratio $M_{2,3}/M_1$, dictating the size of λ_3, is further constrained by leptogenesis: since $N_{2,3}$ have anomalously large Yukawa couplings, on-shell $N_{2,3}$ exchange gives a significant extra washout of n_B, unless $M_{2,3}/M_1$ is large enough that these effects are strongly Boltzmann-suppressed during N_1 decays at $T \sim M_1$. Roughly, one needs $e^{-M_{2,3}/M_1} \lesssim \tilde{m}_1/\tilde{m}_{2,3}$ i.e. $M_{2,3} \gtrsim 10M_1$.

Adopting the computation of [15], this issue is precisely studied in fig. 2b, where we show the maximal n_B/n_γ as a function of λ_3 and of M_2/M_1 for $M_1 = 10^6\,\mathrm{GeV}$, $\tilde{m}_1 = 10^{-3}\,\mathrm{eV}$ and $M_3/M_2 = 2$ (which maximizes $|\varepsilon_1|$, see eq. (26)). We find that for $M_{2,3}/M_1 > 10$ the extra washout effects are negligible, giving λ_3 in agreement with the estimate (27). At smaller $M_{2,3}/M_1$ the CP asymmetry ε_1 becomes larger, but the baryon asymmetry n_B decreases due to $N_{2,3}$-induced washout. One needs $\lambda_3 > 0.1$, which implies $\mathrm{BR}(\mu \to e\gamma)$ close to present experimental limits. This is a robust conclusion, not limited to the specific values of M_1 and of $\tilde{m}_1$ chosen in fig. 2. Indeed the above estimates dictate how fig. 2 can be rescaled to other values of M_1 and $\tilde{m}_1$ (as long as N_1 does not decay far from thermal equilibrium). The precise value of the reheating temperature does not affect n_B, as long as it is a few times larger than M_1. If $T_{\mathrm{RH}} > M_{2,3}$, also $N_{2,3}$ decays contribute to leptogenesis, but without giving significant corrections since they have negligible CP-asymmetries and big washouts.

5 Effects of sub-leading terms

In the previous sections we obtained predictions under the assumption that N_3 has large Yukawa couplings, see eq. (8), which do not give rise to too large neutrino masses thanks to exact texture zeroes, see eq. (9). This patten allows a natural implementation of the mechanism proposed in [20]. A more generic class of natural models is obtained by replacing the texture zeroes with generic entries, as long as they are sufficiently small. Using our vector notation we can now show, without any new computation, that our predictions remain valid also in this case.

A large Yukawa coupling λ_3 is generically needed to lower T_{RH} by enhancing ε_1. Its minimal value can be better determined once sparticles are discovered and their masses measured and understood. The basic assumption that allowed us to control the flavour content $\hat{\lambda}_3$ was that the 32 entry of $\boldsymbol{M}^{-1}$ gives the dominant contribution to seesaw-induced neutrino masses. This assumption followed from a model that naturally accommodates a Yukawa coupling much larger than what is naïvely suggested by the seesaw.

The 33 and 31 entries of $\boldsymbol{M}^{-1}$ would give extra contributions involving the big Yukawa coupling λ_3, if the assumed texture zeros were not exact. But even allowing these extra terms, the neutrino mass matrix can still be written in the form of eq. (14), with $\vec{\lambda}_2$ replaced by a linear combination of $\vec{\lambda}_{1,2,3}$ and with $\vec{\lambda}_3$ unchanged. Therefore, including these extra terms does not affect the prediction for $\hat{\lambda}_3$ of eq. (21) and the consequent prediction for LFV rates.

The 11 and 12 entries of $\boldsymbol{M}^{-1}$ do not give contributions to neutrino masses which involve the large $\vec{\lambda}_3$ Yukawa coupling, and therefore can be neglected as long as λ_1 and λ_2

are sufficiently small. The smallness of λ_2 is demanded by eq. (15) (barring fine-tunings), while the smallness of λ_1 is only suggested by leptogenesis, eq. (13). Predictions for LFV rates can be changed at the expense of a larger $\tilde{m}_1 \gtrsim m_{\rm sun}$, which makes leptogenesis less efficient. This can be compensated by a larger ε_1 obtained from a larger λ_3. In this way (and without caring of the naturalness of the model) it is also possible to suppress LFV effects by concentrating the big Yukawa coupling $\vec{\lambda}_3$ into a single flavour $\ell = e, \mu$ or τ. The big Yukawa coupling λ_3^ℓ still gives detectable effects, by making $\tilde{\ell}_R$ lighter than the two other right-handed sleptons.

6 Conclusions

As pointed out in [20], the DI bound [17, 21] can be evaded, allowing to reconcile thermal leptogenesis with gravitino cosmology within minimal SUSY seesaw leptogenesis. However, the mechanism proposed in [20] employed neutrino Yukawa couplings much larger than suggested by the seesaw mechanism, such that the observed neutrino masses were reproduced thanks to cancellations.

In this work we sketched characteristics of models, which achieve this goal avoiding accidental cancellations by employing the right-handed neutrino mass matrix of the form (9) and large Yukawa couplings $N_3(\vec{\lambda}_3 \cdot \vec{L})H$ with $\lambda_3 \sim 0.1$. The texture (9) allows to control the contribution of the large neutrino Yukawa couplings in the seesaw mechanism avoiding contribution to neutrino masses much larger than their observed values. We outlined how flavour models can justify the texture zeroes in eq. (9). Assuming also $\tilde{m}_1 \sim 10^{-3}\,\text{eV} \ll m_{\rm sun}$ (so that thermal leptogenesis is maximally efficient), we found that the flavour structure of the big Yukawa coupling is related to neutrino masses and mixings (see eq. (21)). In conclusion, the models predict a detectable $\text{BR}(\mu \to e\gamma)$ and a precise correlation with $\text{BR}(\tau \to e\gamma)$ and $\text{BR}(\tau \to \mu\gamma)$.

We would like to finish by remarking that, although we do not claim that our models are the only ones that achieve the goal described above, we could not find any other solution. Similarly, although the smallness of $\tilde{m}_1$ is not implied by the previous assumptions, in most of our parameter space it is difficult to avoid it. Future measurements of sparticle masses can clarify this issue. In this weak sense the predictions we obtained might be a generic testable consequence of natural realizations of the mechanism suggested in [20].

Acknowledgments We thank S. Davidson for many discussions. A.S. thanks F. Feruglio. K.T. is indebted to S. Pokorski for continuous support and to P. H. Chankowski for useful comments. K.T. also thanks the European Centre for Theoretical Studies in Nuclear Physics and Related Areas in Villazzano (Trento), Italy, where he participated in the *Marie Curie* Doctoral Training Programme, for hospitality and stimulating atmosphere during the preparation of this work. The work of M. R. is partially supported by the ESF Grant 6140, by the EC MC contract MERG-CT-2003-503626 and by the Ministry of Education and Research of the Republic of Estonia. The work of K. T. is partially supported by the Polish State Committee for Scientific Research Grants 2 P03B 129 24 for years 2003-05 and 1 P03D 014 26 for years 2004-06.

References

[1] P. Minkowski, Phys. Lett. B 67 (1977) 421. See also M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, edited by P. van Nieuwenhuizen and D. Freedman, (North-Holland, 1979), p. 315; S.L. Glashow, in Quarks and Leptons, Cargése, eds. M. Lévy et al., (Plenum, 1980, New-York), p. 707; T. Yanagida, in Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe, edited by O. Sawada and A. Sugamoto (KEK Report No. 79- 18, Tsukuba, 1979), p. 95; R.N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44 (1980) 912.

[2] M. Fukugita and T. Yanagida, Phys. Lett. B174 (1986) 45.

[3] For recent analyses see R.H. Cyburt, J. Ellis, B.D. Fields, K.A. Olive, Phys. Rev. D67 (2003) 103521; M. Kawasaki, K. Kohri, T. Moroi, astro-ph/0402490.

[4] M. Bolz, A.Brandenburg, W. Buchmüller, Nucl. Phys. B606 (2003) 518.

[5] M. Flanz, E.A. Paschos and U. Sarkar, Phys. Lett. B345 (1995) 248; M. Flanz, E.A. Paschos, U. Sarkar and J. Weiss, Phys. Lett. B389 (1996) 693.

[6] A. Pilaftsis, Phys. Rev. D56 (1997) 5431; Int. J. Mod. Phys. A17 (1999) 575.

[7] J. R. Ellis, M. Raidal and T. Yanagida, Phys. Lett. B 546 (2002) 228.

[8] A. Pilaftsis, T.E.J. Underwood, Nucl. Phys. B692 (2004) 303.

[9] K. Turzynski, Phys. Lett. B589 (2004) 135.

[10] Y. Grossman, T. Kashti, Y. Nir, E. Roulet, Phys. Rev. Lett. 91 (2003) 251801 and hep-ph/0407063; G. D'Ambrosio, G.F. Giudice, M. Raidal, Phys. Lett. B575 (2003) 75.

[11] E.J. Chun, Phys. Rev. D69 (2004) 117303; T. Hambye, J. March-Russell, S.M. West, JHEP 0407 (2004) 070; L. Boubekeur, T. Hambye, G. Senjanovic, Phys. Rev. Lett. 93 (2004) 111601 [hep-ph/0404038].

[12] G. Lazarides, Q. Shafi, Phys. Lett. B258 (1991) 305; R. Allahverdi, B. Dutta, A. Mazumdar, Phys. Rev. D67 (2003) 123515; R. Allahverdi, M. Drees, Phys. Rev. D69 (2004) 103522.

[13] K. Hamaguchi, H. Murayama, T. Yanagida, Phys. Rev. D65 (2002) 043512.

[14] J. R. Ellis, M. Raidal and T. Yanagida, Phys. Lett. B 581 (2004) 9.

[15] G.F. Giudice, A. Notari, M. Raidal, A. Riotto, A. Strumia, Nucl. Phys. B685 (2004) 89.

[16] H.S. Goh, M.A. Luty, S.P. Ng, Phys. Rev. D69 (2004) 014002 [hep-th/0309103] and references therein.

[17] S. Davidson, A. Ibarra, Phys. Lett. B535 (2002) 25. The bound (5) on ε_1 was also present in [13]. An earlier work ([21] around eq. (5.1)) presented this bound as valid only up to cancellations, in agreement with our present findings.

[18] For a recent review see W. Buchmüller, P. Di Bari, M. Plümacher, New J. Phys. 6 (2004) 105. We do not fully agree with some comments. Concerning 'gauge scatterings', we are unaware of any contradiction between [15] and the final version of [8]. Anyhow, an independent computation would be opportune, since only [15] attempted to include precisely these effects, plagued by tricky IR divergences. Furthermore, related effects (3 body decays, radiative corrections to $N_1 \to LH$ decay and its asymmetry) remain to be computed. As for the 'spectator processes', we think that this is another way of describing the flavour factors, already studied in [21] with temperature dependence of the sphaleron rate taken into account (see also paragraph 'Sphalerons' in [15]).

[19] L. Covi, E. Roulet, F. Vissani, Phys. Lett. B384 (1996) 169.

[20] T. Hambye, Y. Lin, A. Notari, M. Papucci, A. Strumia, Nucl. Phys. B695 (2004) 169; S. Davidson and R. Kitano, JHEP 043 (2004) 020.

[21] R. Barbieri, P. Creminelli, A. Strumia, N. Tetradis, Nucl. Phys. B575 (2000) 61.

[22] See e.g. K.S. Babu and S.M. Barr, Phys. Lett. B381 (1996) 202; J. Sato and T. Yanagida, Phys. Lett. B493 (2000) 356.

[23] J. K. Elwood, N. Irges and P. Ramond, Phys. Rev. Lett. 81 (1998) 5064 and section 9.1.2 of the review by G. Altarelli, F. Feruglio, New J. Phys. 6 (2004) 106 [hep-ph/0405048].

[24] See e.g. F. Feruglio, A. Strumia, F. Vissani, Nucl. Phys. B637 (2002) 345 [hep-ph/0201291]; S.F. King, Phys. Rev. D67 (2003) 113010; V. Barger, D.A. Dicus, H.-J. He, T.-J. Li, Phys. Lett. B583 (2004) 173; A. Ibarra, G.G. Ross, Phys. Lett. B591 (2004) 285 [hep-ph/0312138]; J.-W. Mei, Z.-Z. Xing, Phys. Rev. D69 (2004) 073003.

[25] P.H. Chankowski, J.R. Ellis, S. Pokorski, M. Raidal, K. Turzynski, Nucl. Phys. B690 (2004) 279.

[26] CHOOZ collaboration, Eur. Phys. J. C27 (2003) 331.

[27] See e.g. P. H. Frampton, S. L. Glashow and T. Yanagida, Phys. Lett. B **548** (2002) 119 [hep-ph/0208157]; M. Raidal and A. Strumia, Phys. Lett. B **553** (2003) 72 [hep-ph/0210021]; S. Raby, Phys. Lett. B **561** (2003) 119 [hep-ph/0302027].

[28] J. A. Casas and A. Ibarra, Nucl. Phys. B **618** (2001) 171 [hep-ph/0103065].

[29] P.H. Chankowski, S. Pokorski, Int. J. Mod. Phys. A17 (2002) 575.

[30] J. Hisano, D. Nomura, Phys. Rev. D59 (1999) 116005.

Addendum to
Low-scale standard supersymmetric leptogenesis

Martti Raidal[1], **Alessandro Strumia**[2] and **Krzysztof Turzyński**[3]

[1] *National Institute of Chemical Physics and Biophysics, Tallinn 10143, Estonia*
[2] *Dipartimento di Fisica dell'Università di Pisa and INFN, Italia*
[3] *Institute of Theoretical Physics, Warsaw University, ul. Hoża 69, 00-681 Warsaw, Poland*

In [1A] we proposed a supersymmetric leptogenesis model where the two heavier right-handed neutrinos $N_{2,3}$ have Yukawa couplings naturally larger than what suggested by neutrino masses. This allows to avoid the potential conflict between supersymmetry and thermal leptogenesis [2A] by enhancing the CP-asymmetry ε_1 in N_1 decays without significantly reducing the efficiency η.

As pointed to us by S. Davidson, when studying the reduction in η we only considered the wash-out produced by on-shell exchange of $N_{2,3}$: off-shell exchange of $N_{2,3}$ can give an extra important contribution.

It is useful to understand from a more general point of view that off-shell wash-out is directly linked to the mechanism that enhances ε_1. While in the standard leptogenesis scenario ε_1 is dominantly generated by the dimension-5 neutrino mass operator mediated by $N_{2,3}$, in our scenario $N_{2,3}$ mediate the dimension-7 operator $(LH)\partial^2(LH)$, that dominantly contributes both to ε_1 and to off-shell wash-out. It is therefore crucial to ensure that a sizable enhancement of ε_1 does not imply a sizable reduction of η. The relevant non-renormalizable effective operators present at energies around the N_1 mass are the supersymmetrized version of

$$\mathscr{L}_{\text{eff}} = c_5\frac{(LH)^2}{2} + c_7\frac{(LH)\partial^2(LH)}{2} + \text{h.c.} \tag{A.1}$$

In our model $c_5 = 0$ and $c_7 = -\lambda_3^2(M_3 - M_2)/M_2^2 M_3^2$. Neglecting the $c_5 \cdot c_7$ interference term, these operators contribute to the equilibrium space-time densities of $\Delta L = 2$ wash-out scatterings at temperature T as

$$\gamma_{Ns}(LH \leftrightarrow \bar{L}\bar{H}) = \gamma_{Nt}(LL \to \bar{H}\bar{H}) \overset{T\ll M_{2,3}}{\simeq} \frac{3T^6}{8\pi^5}|c_5|^2 + \frac{120T^{10}}{\pi^5}|c_7|^2. \tag{A.2}$$

We followed the notation of [3A]. Analogous expressions hold for $\Delta L = 2$ scatterings involving sparticles. The CP asymmetry is

$$\varepsilon_1 = \frac{3M_1}{8\pi}\text{Im}\, c_5 - \frac{3M_1^3}{16\pi}\text{Im}\, c_7 \tag{A.3}$$

in agreement with eq. (26) of [1A]. The above operator approximation has a general validity and shows that a too large off-shell washout would exclude the whole scenario, not only our model. One can analytically check that in most of the parameter space off-shell washout is small enough.

We here present a full numerical computation of thermal leptogenesis that includes all contributions to wash-out as predicted by our supersymmetric model. Notice that the generic



Figure 3: *Fig. 2b of* [1A] *modified taking into account all contributions to* $\Delta L = 2$ *scatterings. We assumed* $M_1 = 10^6\,\mathrm{GeV}$, $M_3/M_2 = 2$, $\tilde{m}_1 = 10^{-3}\,\mathrm{eV}$.

expressions for the $\Delta L = 2$ wash-out scattering rates [4A] are not sufficiently accurate in our very specific context. Indeed these expressions are obtained by modifying the propagators of the mass-eigenstate right-handed neutrinos N_i and sneutrinos $\tilde{N}_i$ ($i = \{1,2,3\}$) by taking into account the decay widths Γ_i according to the usual prescription:

$$\Pi_{ii}(p) = \frac{1}{p^2 - M_i^2 + i\Gamma_i M_i}. \tag{A.4}$$

We see that $\Pi_{ii}(0)$ gets modified, spoiling the delicate cancellation that in our model suppresses the dimension-5 operator. To correctly reproduce this cancellation one has to take into account that also the off-diagonal Π_{ij} propagators develop an imaginary part.

Fig. 3 shows our full result, that can be directly compared with fig. 2b in the paper [1A], where the off-shell effect was not included. The new off-shell washout is more important than on-shell washout only when $\lambda_3 \sim M_2/M_1 \sim 1$. As a result, the region where leptogenesis can be successful survives and is shaded in green in fig. 3, getting somewhat reduced. The constraint on the parameter that controls lepton-flavour violating signals remains unaltered: $\lambda_3 \gtrsim 0.1$.

References

[1A] M. Raidal, A. Strumia and K. Turzynski, Phys. Lett. B 609 (2005) 351 [hep-ph/0408015].

[2A] S. Davidson, A. Ibarra, Phys. Lett. B535 (2002) 25.

[3A] G.F. Giudice, A. Notari, M. Raidal, A. Riotto, A. Strumia, Nucl. Phys. B685 (2004) 89.

[4A] M. Plumacher, Nucl. Phys. B 530 (1998) 207 [hep-ph/9704231].